

## Fungicide for Control of Foliar Diseases of Small Grains

Fungicide	Product	Rust <sup>a</sup>	Powdery Mildew	Septoria Leaf Blotch	Glume Blotch	Crop <sup>b</sup>	Days to Harvest	Remarks and suggestions
azoxystrobin	Quadris	X	X	X	X	W		Apply at flag-leaf stage before ligule of flag leaf emerges. Do not graze or use for feed. May be used for bedding. Also labeled for use on other diseases.
mancozeb	Dithane M-45	X		X	X	B / O / R / W	26	Don't graze within 26 days after application. Maximum of three applications per crop season.
	Manzate 200	X		X	X	B / O / R / W	26	
	Penncozeb	X		X	X	B / O / R / W	26	
triadimefon	Bayleton <sup>c</sup>	X	X	X		B/ W	21	Also labeled for use on barley scald. Do not exceed a total of 16 oz per crop season.
propiconazole	Tilt	X	X	X	X	B/ R/ W		Apply at flag-leaf stage before ligule of flag leaf emerges. Only one application per crop season.
pyraclostrobin	Headline	X	X	X	X	B/ R/ W		
propiconazole+	Stratego	X	X	X	X	B/ R/ W		
trifloxystrobin triadimefon	Baytan 30	X	X			B/ R/ W		Do not graze or use for feed. May be used for bedding. Also labeled for use on other diseases.

<sup>a</sup>Rust includes stem, leaf, and stripe rust fungi.

<sup>b</sup>Corn abbreviations: B = barley, O = oat, R = rye, W = wheat

<sup>c</sup>Bayleton can be tank-mixed with formulations of mancozeb.